

Chemical Kinetics

❖ Rate of Reaction:

The quantity of a reactant species consumed or the quantity of a product species formed in unit time in a chemical reaction is called the rate of the reaction.


Consider a reaction, $A \rightarrow B$

$$\text{Rate} = -\frac{d[A]}{dt} = +\frac{d[B]}{dt}$$

–ve sign indicates that with progress of time concentration of A will decrease.

❖ Rate Laws & Rate Constant:

Let us consider a general reaction,



According to rate law,

$$\text{Rate} \propto [A][B]$$

$$\text{Or, Rate} = K[A][B]$$

This is known as rate law & ' K ' is called rate constant which is independent of concentration but depend on temperature.

➤ Mention the factors affect the rate of a chemical reaction.


Factors affecting the rate of a chemical reaction are following –

- i. Nature of reactants
- ii. Concentration of reactants
- iii. Effect of temperature
- iv. Surface area of reactants
- v. Presence of catalyst
- vi. Presence of light

❖ Order of a Reaction:

Order of a reaction is defined as the sum of the powers or exponents to which the molar concentrations are raised in the rate law or rate equation.

Let us consider a general reaction,




According to rate law,

$$R = K[A]^a[B]^b$$

where a & b may or may not be equal to l & m respectively. Here the overall order of the reaction is $(a + b)$.


- ✓ Order of a reaction is purely experimental quantity.
- ✓ Order of reaction may be fraction, zero or even negative. As for example, thermal decomposition of acetaldehyde,



has order equal to $3/2$.

❖ Example of Reactions having Different Order:



- I. Zero order reaction: Dissociation of ammonia on tungsten or dissociation of HI on gold surface.
- II. First order reaction: Conversion of cyclopropane to propylene, substitution reaction of tertiary alkyl halide by nucleophile.
- III. Second order reaction: $H_2 + I_2 \rightarrow 2HI$, substitution reaction of primary alkyl halide by nucleophile.
- IV. Fractional order reaction: Thermal decomposition of acetaldehyde,



has order equal to $3/2$.

❖ Elementary Reaction:

A chemical reaction that takes place in one & only one step, i.e., all that occurs in a single step is called elementary reaction. For example –



Molecularity of an elementary reaction is the number of the reactant particles that are involved in each individual chemical event.

❖ Molecularity of Elementary Reaction:

It is defined as the number of reactant particles (atoms, molecules, free radicals or ions) that are involved in each individual chemical event.

If a reaction takes place in a number of steps then slowest step is usually the rate determining step. Molecularity of the rate determining step is said to be the molecularity of the reaction.

- ✓ Molecularity of a reaction is purely theoretical concept.
- ✓ Molecularity of a reaction is necessarily a positive integer.

➤ Elementary reactions with molecularity greater than three generally do not occur – justify.


The reactants are activated as a result of molecular collisions. But the collisions generally require high amount of energy because of inter – nuclear & inter – electronic repulsions. For a collision minimum two molecules are must. If the molecularity of an elementary step is 3, then three molecules must collide requiring high amount of energy. For a four – body collision, energy required will be still higher & therefore unknown. It is far easier to have two – body collisions & that will take place easily. Therefore, elementary reaction of molecularity 4 or greater is unknown. If molecularity becomes greater than 3, then such reaction involve two or more steps, such that each step has its own molecularity not greater than 3.

❖ Difference Between Molecularity & Order:

The main differences are –

- I. Order is an experimental quantity whereas molecularity of a reaction is a purely theoretical concept.
- II. Order of a reaction is not necessarily integer. It may be fraction, zero or even negative. Whereas molecularity of a reaction is necessarily a positive integer.
- III. Order may change with physical condition, such as pressure, solvent etc whereas molecularity is an invariant number for a given chemical reaction.


➤ Give an example of a chemical reaction where molecularity & order is different.



So, rate = $k[\text{CH}_3\text{CHO}]^{3/2}$. Here the order of the reaction is 1.5 but molecularity is 1. Hence order & molecularity is different.

❖ Integrated Rate Laws:

1. **Zero Order Reaction:** A reaction whose rate is independent of concentration is called a zero order reaction. Thus for a zero order reaction –



$$\text{So, rate} = -\frac{d[A]}{dt} = k[A]^0 = k$$

where $[A]$ is the concentration of the reactant A at time t , thus –

$$\int_a^{[A]} d[A] = -k \int_0^t dt$$

$$\text{Or, } [A] - a = -kt$$


$$\text{Or, } [A] = a - kt$$

A zero order reaction will be completed when $[A] = 0$, so,

$$a - kt_{\text{completion}} = 0$$

$$\text{Or, } t_{\text{completion}} = \left(\frac{a}{k}\right)$$

Graphical Representation: If we plot $[A]$ vs t , we get a straight line with slope = $-k$.



Half – life period: When $t = t_{\frac{1}{2}}$, then, $[A] = a/2$, now –

$$\frac{a}{2} = a - kt_{\frac{1}{2}}$$

$$\text{Or, } t_{\frac{1}{2}} = \left(\frac{a}{2k}\right), \text{ i. e., } t_{1/2} \propto a$$

Unit of rate constant: $\text{mol. L}^{-1}\text{sec}^{-1}$

Characteristics:

- I. Rate is independent of instantaneous concentration of the reactant.
- II. Unit of rate & rate constant are the same.
- III. Zero order reaction must be multistep in nature.
- IV. Zero order reaction can be completed in a finite time.
- V. Half – life period is directly proportional to initial concentration of reactant.

➤ Write down the unit of the rate constant of a zero order reaction.

$$\text{mol. L}^{-1}\text{sec}^{-1}$$

➤ What do you mean by half-life period of a reaction?

The half-life of a reaction, $t_{1/2}$, is the amount of time needed for a reactant concentration to decrease by half compared to its initial concentration.

➤ Zero order reaction is multistep — Explain.

If zero order reaction is single step then the reaction would be an elementary reaction & in that case order of the reaction would be same as molecularity. But molecularity of a reaction cannot be zero. So zero order reaction must be a multistep reaction. The reactant with respect to which the order becomes zero is involved in a step which is not the rate determining step (r.d.s.).

➤ Does a zero order reaction go to completion?

Rate equation for a zero order reaction —

$$-\frac{dc}{dt} = kC^0$$

$$\text{Or, } \int dc = - \int k dt$$

$$\text{Or, } C = -kt + IC$$

$$\text{at } t = 0, C = C_0, \text{ then } IC = C_0$$

$$\text{So, } C = C_0 - kt$$

Hence, a zero order reaction will go to completion & time duration is $\frac{C_0}{k}$.

➤ For a reaction the plot of $\log t_{1/2}$ versus $\log a$ turns out to be a straight line with positive slope and intercept, which makes an angle of 45° with the $\log a$ axis. What is the order of the reaction?

We know half – life period of a nth order reaction is given as –

$$t_{1/2} = \frac{k}{a^{n-1}}$$

$$\text{Or, } \log t_{1/2} = \log k + (1 - n) \log a$$

Slope of $\log t_{1/2}$ vs $\log a = (1 - n)$.

So we can write –

$$(1 - n) = \tan 45^\circ = 1, \quad \text{Or,} \quad n = 0$$

Hence the given reaction is a zero order reaction.

- In a certain kinetic study, the plot of $\log t_{1/2}$ vs $\log a$ was found to be a straight line with unit slope. State with reason the order of the reaction.

We know half – life period of a nth order reaction is given as –

$$t_{1/2} = \frac{k}{a^{n-1}}$$

$$\text{Or, } \log t_{1/2} = \log k + (1 - n) \log a$$

Slope of $\log t_{1/2}$ vs $\log a = (1 - n)$.


So we can write –

$$(1 - n) = 1$$

$$\text{Or, } n = 0$$

Hence the given reaction is a zero order reaction.

2. **First Order Reaction:** A first order reaction is one whose rate is linearly proportional to the instantaneous concentration of the reactant. Let us consider a first order reaction –



Let initial concentration of reactant is $[A]_0 = a$ & at an arbitrary time t it is $[A] = a - x$. Then the differential rate law is given as –

$$-\frac{d(a - x)}{dt} = k(a - x)$$

$$\text{Or, } \left(\frac{dx}{dt}\right) = k(a - x)$$


$$\text{Or, } \int_0^x \frac{dx}{(a - x)} = k \int_0^t dt$$

$$\text{Or, } \ln \left(\frac{a}{a - x}\right) = kt$$

$$\text{Or, } \log \left(\frac{a}{a - x}\right) = \frac{k}{2.303} t$$

$$\text{Or, } t = \frac{2.303}{k} \log \left(\frac{a}{a - x}\right)$$

Graphical Representation: If we plot $\log \left(\frac{a}{a - x}\right)$ vs t , we get a straight line passing through the origin & slope = $k/2.303$.



Half – life period: When $t = t_{\frac{1}{2}}$, then, $x = a/2$, from above equation –

$$t_{\frac{1}{2}} = \frac{2.303}{k} \log \left(\frac{a}{a - \frac{a}{2}} \right) = \frac{1}{k} \ln 2 = \frac{0.693}{k}$$

So, half – life period is independent of initial concentration of reactant.

Unit of rate constant: sec^{-1}

Characteristics:

- I. First order reaction never goes to completion.
- II. Half – life period is independent of initial concentration of reactant.

➤ First order reaction never proceeds to completion – justify or criticize.

For first order reaction –

$$\frac{[A]}{[A]_0} = e^{-kt}$$

When reaction will be completed then $[A]$ should be equal to zero & it will be possible only when $t = \infty$, so first order reaction never goes to completion.

➤ Show for a first order reaction, the time required for 99.9% completion of the reaction is almost 10 times the time for 50% completion.

For a first order reaction,

$$t = \frac{1}{k} \ln \frac{a}{a-x} = \frac{2.303}{k} \log \frac{a}{a-0.999a}$$

$$\text{Or, } t = \frac{2.303}{k} \log 10^3 = \frac{6.909}{k}$$

$$\text{We know, } t_{\frac{1}{2}} = \frac{0.693}{k}$$

$$\text{So, } \frac{t}{t_{\frac{1}{2}}} = \frac{6.909}{0.693} \cong 10.$$

Hence, $t = 10 \times t_{1/2}$ (proved)

➤ Show that the time required for 75% completion of a first – order reaction is twice the time required for 50% completion of the reaction.

For a first order reaction,

$$t = \frac{1}{k} \ln \frac{a}{a-x} = \frac{2.303}{k} \log \frac{a}{a-0.75a}$$


$$\text{Or, } t = \frac{2.303}{k} \log 4 = \frac{1.3865}{k}$$

$$\text{We know, } t_{\frac{1}{2}} = \frac{0.693}{k}$$

$$\text{So, } \frac{t}{t_{\frac{1}{2}}} = \frac{1.3865}{0.693} \cong 2.$$

Hence, $t = 2 \times t_{1/2}$ (proved)

3. **Second Order Reaction:** Consider a second order reaction involving only one reactant, i.e.



Let initial concentration of reactant is $[A]_0 = a$ & at an arbitrary time t it is $[A] = a - x$. Then the differential rate law is given as –

$$-\frac{d(a-x)}{dt} = k(a-x)^2$$

$$\text{Or, } \left(\frac{dx}{dt}\right) = k(a-x)^2$$

$$\text{Or, } \int_0^x \frac{dx}{(a-x)^2} = k \int_0^t k dt$$


$$\text{Or, } \left(\frac{1}{a-x}\right) - \frac{1}{a} = kt$$

$$\text{Or, } \frac{1}{a-x} = \frac{1}{a} + kt$$

$$\text{Or, } kt = \frac{x}{a(a-x)}$$

$$\text{Or, } t = \frac{1}{k} \times \frac{x}{a(a-x)}$$

Graphical Representation: If we plot $1/[R]$ vs t , we get a straight line with intercept $= 1/a$ & slope = k .



Half – life period: When $t = t_{\frac{1}{2}}$, then, $x = a/2$, from above equation –

Half – life period: When $t = t_{\frac{1}{2}}$, then, $x = a/2$, from above equation –

$$t_{\frac{1}{2}} = \frac{1}{k} \times \frac{\frac{a}{2}}{a \cdot \frac{a}{2}} = \frac{1}{ak}$$

Hence, half – life period is inversely proportional to initial concentration of reactant.

Unit of rate constant: $\text{mol}^{-1}\text{lit. sec}^{-1}$

Characteristics:

- I. Second order reaction transfers to first order reaction when one of the reactant is present in large excess.
- II. Half – life period of second order reaction is inversely proportional to initial concentration of reactant.

- Write down the unit of rate constant of a 2nd order reaction.

$$\text{mol}^{-1}\text{lit. sec}^{-1}$$

- Second order reaction never proceeds to completion – justify or criticize.

$$\text{For a 2nd order reaction, } k = \frac{1}{t} \frac{x}{a(a-x)}$$

$$\text{So, } t = \frac{1}{k} \frac{x}{a(a-x)}$$

When $t = t_{\text{completion}}$, then $a = 0$,

$$\text{So, } t_{\text{completion}} = \frac{1}{k} \cdot \frac{x}{0} = \infty$$

So, 2nd order reaction never goes to completion.

- For a second order reaction $A \rightarrow \text{products}$. Hence show that the time required for three – fourths of the reactant to decay ($t_{3/4}$) is equal to $3t_{1/2}$.

For a second order reaction –

$$\frac{1}{a-x} = \frac{1}{a} + kt$$

$$\text{Or, } \frac{1}{a - \frac{3a}{4}} = \frac{1}{a} + kt_{\frac{3}{4}} \quad (\text{for } \frac{3}{4} \text{ th of the reactant})$$

$$\text{Or, } \frac{1}{\frac{a}{4}} = \frac{1}{a} + kt_{\frac{3}{4}}$$

$$\text{Or, } kt_{\frac{3}{4}} = \frac{4}{a} - \frac{1}{a} = \frac{3}{a}$$

$$\text{Or, } t_{\frac{3}{4}} = 3 \times \frac{1}{ka} = 3t_{1/2} \quad (\text{we know for 2nd order reaction } t_{1/2} = \frac{1}{ka})$$

$$\text{Or, } t_{3/4} = 3 \times t_{1/2} \quad (\text{proved})$$